

Exercise sheet № 2 - Introduction, symmetries

(Suggestion: draw the Feynman diagrams of all the reactions.)

1. What is the relative angular momentum ℓ in the final state of the decay $\Delta^{++} \rightarrow p\pi^+$?
2. Find the proportion of the two decays $K^{*+} \rightarrow K^0\pi^+$ and $K^{*+} \rightarrow K^+\pi^0$. For each of them, obtain a decay with a similar branching fraction by a simple rotation in the isospin space.
3. Find the relative proportions of the decays $\Sigma^{*0} \rightarrow \Sigma^+\pi^-$, $\Sigma^{*0} \rightarrow \Sigma^0\pi^0$ and $\Sigma^{*0} \rightarrow \Sigma^-\pi^+$.
4. The largest branching fraction of ρ^0 decays corresponds to $\rho^0 \rightarrow \pi\pi$.
 - (a) Which interaction is responsible for this decay? Give as much arguments as possible.
 - (b) Examine the conservation of P in the decay. Using symmetry considerations, show that the final state $\pi^0\pi^0$ is forbidden.
 - (c) Study the conservation of C in the decay. What can be concluded?
 - (d) Draw the same conclusion by looking at isospin conservation.
 - (e) Draw the Feynman diagram of the decay.
 - (f) Interpret the branching fraction of the decay $\rho^0 \rightarrow \pi^+\pi^-\pi^0$ and that of $\rho^0 \rightarrow \pi^+\pi^-\pi^0\pi^0$.
5. Study the coherence between the different quantum numbers (J^{PC} from the PDG, orbital angular momentum from the particle name) of the $c\bar{c}$ mesons: $\eta_c(1S)$, $J/\psi(1S)$, $\chi_{c0}(1P)$, $h_c(1P)$ and $\psi(2S)$. For each meson, give the possible pure spin state(s).
6. Using the relevant discrete symmetry (and the right arguments), show that the decay $\omega \rightarrow \pi^0\gamma$ is allowed, but that $\omega \rightarrow \pi^0\gamma\gamma$ is forbidden.
7. Show that the two- and three-pions final states in the decays of neutral kaons are well defined CP eigenstates. Find their CP contents.
8. The meson $\phi(1020)$ is a $s\bar{s}$ state, decaying to K^+K^- and $K^0\bar{K}^0$ (through which interaction?).
 - (a) Study the conservation of P and C in these decays.
 - (b) Making the (good) approximation that K_S^0 and K_L^0 are CP eigenstates with $CP = +1$ and $CP = -1$, respectively, explain the fact that the $K^0\bar{K}^0$ final state is observed only as $K_S^0K_L^0$, and not as $K_S^0K_S^0$ or $K_L^0K_L^0$.
9. Obtain a general expression for the kinematic boundaries of the Dalitz plot. Interpret the corners of the allowed region in term of momenta of the final state particles. Apply to the decay $B^0 \rightarrow D^0K^+\pi^-$.